

1 negbinom: Negative Binomial Regression for Event Count Dependent Variables

Use the negative binomial regression if you have a count of events for each observation of your dependent variable. The negative binomial model is frequently used to estimate over-dispersed event count models.

1.0.1 Syntax

```
> z.out <- zelig(Y ~ X1 + X2, model = "negbinom", data = mydata)
> x.out <- setx(z.out)
> s.out <- sim(z.out, x = x.out)
```

1.0.2 Additional Inputs

In addition to the standard inputs, `zelig()` takes the following additional options for negative binomial regression:

- **robust**: defaults to `FALSE`. If `TRUE` is selected, `zelig()` computes robust standard errors via the `sandwich` package (see [7]). The default type of robust standard error is heteroskedastic and autocorrelation consistent (HAC), and assumes that observations are ordered by time index.

In addition, **robust** may be a list with the following options:

- **method**: Choose from
 - * `"vcovHAC"`: (default if **robust** = `TRUE`) HAC standard errors.
 - * `"kernHAC"`: HAC standard errors using the weights given in [1].
 - * `"weave"`: HAC standard errors using the weights given in [3].
- **order.by**: defaults to `NULL` (the observations are chronologically ordered as in the original data). Optionally, you may specify a vector of weights (either as **order.by** = `z`, where `z` exists outside the data frame; or as **order.by** = `~z`, where `z` is a variable in the data frame). The observations are chronologically ordered by the size of `z`.
- **...**: additional options passed to the functions specified in **method**. See the `sandwich` library and [7] for more options.

1.0.3 Example

Load sample data:

```
> data(sanction)
```

Estimate the model:

```
> z.out <- zelig(num ~ target + coop, model = "negbinom", data = sanction)
> summary(z.out)
```

Set values for the explanatory variables to their default mean values:


```
> x.out <- setx(z.out)
```

Simulate fitted values:

```
> s.out <- sim(z.out, x = x.out)
```

```
> summary(s.out)
```

```
> plot(s.out)
```



1.0.4 Model

Let Y_i be the number of independent events that occur during a fixed time period. This variable can take any non-negative integer value.

- The negative binomial distribution is derived by letting the mean of the Poisson distribution vary according to a fixed parameter ζ given by the Gamma distribution. The *stochastic component* is given by

$$\begin{aligned} Y_i | \zeta_i &\sim \text{Poisson}(\zeta_i \mu_i), \\ \zeta_i &\sim \frac{1}{\theta} \text{Gamma}(\theta). \end{aligned}$$

The marginal distribution of Y_i is then the negative binomial with mean μ_i and variance $\mu_i + \mu_i^2/\theta$:

$$\begin{aligned} Y_i &\sim \text{NegBinom}(\mu_i, \theta), \\ &= \frac{\Gamma(\theta + y_i)}{y! \Gamma(\theta)} \frac{\mu_i^{y_i} \theta^\theta}{(\mu_i + \theta)^{\theta + y_i}}, \end{aligned}$$

where θ is the systematic parameter of the Gamma distribution modeling ζ_i .

- The *systematic component* is given by

$$\mu_i = \exp(x_i \beta)$$

where x_i is the vector of k explanatory variables and β is the vector of coefficients.

1.0.5 Quantities of Interest

- The expected values (`qi$ev`) are simulations of the mean of the stochastic component. Thus,

$$E(Y) = \mu_i = \exp(x_i \beta),$$

given simulations of β .

- The predicted value (`qi$pr`) drawn from the distribution defined by the set of parameters (μ_i, θ) .
- The first difference (`qi$fd`) is

$$\text{FD} = E(Y|x_1) - E(Y|x)$$

- In conditional prediction models, the average expected treatment effect (`att.ev`) for the treatment group is

$$\frac{1}{\sum_{i=1}^n t_i} \sum_{i:t_i=1}^n \{Y_i(t_i = 1) - E[Y_i(t_i = 0)]\},$$

where t_i is a binary explanatory variable defining the treatment ($t_i = 1$) and control ($t_i = 0$) groups. Variation in the simulations are due to uncertainty in simulating $E[Y_i(t_i = 0)]$, the counterfactual expected value of Y_i for observations in the treatment group, under the assumption that everything stays the same except that the treatment indicator is switched to $t_i = 0$.

- In conditional prediction models, the average predicted treatment effect (`att.pr`) for the treatment group is

$$\frac{1}{\sum_{i=1}^n t_i} \sum_{i:t_i=1}^n \{Y_i(t_i = 1) - \widehat{Y_i(t_i = 0)}\},$$

where t_i is a binary explanatory variable defining the treatment ($t_i = 1$) and control ($t_i = 0$) groups. Variation in the simulations are due to uncertainty in simulating $\widehat{Y_i(t_i = 0)}$, the counterfactual predicted value of Y_i for observations in the treatment group, under the assumption that everything stays the same except that the treatment indicator is switched to $t_i = 0$.

1.0.6 Output Values

The output of each Zelig command contains useful information which you may view. For example, if you run `z.out <- zelig(y ~ x, model = "negbinom", data)`, then you may examine the available information in `z.out` by using `names(z.out)`, see the `coefficients` by using `z.out$coefficients`, and a default summary of information through `summary(z.out)`. Other elements available through the `$` operator are listed below.

- From the `zelig()` output object `z.out`, you may extract:
 - `coefficients`: parameter estimates for the explanatory variables.
 - `theta`: the maximum likelihood estimate for the stochastic parameter θ .
 - `SE.theta`: the standard error for `theta`.
 - `residuals`: the working residuals in the final iteration of the IWLS fit.
 - `fitted.values`: a vector of the fitted values for the systemic component λ .
 - `linear.predictors`: a vector of $x_i\beta$.
 - `aic`: Akaike's Information Criterion (minus twice the maximized log-likelihood plus twice the number of coefficients).
 - `df.residual`: the residual degrees of freedom.
 - `df.null`: the residual degrees of freedom for the null model.
 - `zelig.data`: the input data frame if `save.data = TRUE`.
- From `summary(z.out)`, you may extract:
 - `coefficients`: the parameter estimates with their associated standard errors, p -values, and t -statistics.
 - `cov.scaled`: a $k \times k$ matrix of scaled covariances.
 - `cov.unscaled`: a $k \times k$ matrix of unscaled covariances.
- From the `sim()` output object `s.out`, you may extract quantities of interest arranged as matrices indexed by simulation \times x -observation (for more than one x -observation). Available quantities are:

- `qi$ev`: the simulated expected values given the specified values of `x`.
- `qi$pr`: the simulated predicted values drawn from the distribution defined by (μ_i, θ) .
- `qi$fd`: the simulated first differences in the simulated expected values given the specified values of `x` and `x1`.
- `qi$att.ev`: the simulated average expected treatment effect for the treated from conditional prediction models.
- `qi$att.pr`: the simulated average predicted treatment effect for the treated from conditional prediction models.

How to Cite the Negative Binomial Model

Kosuke Imai, Olivia Lau, and Gary King. *negbinom: Negative Binomial Regression for Event Count Dependent Variables*, 2011

How to Cite the Zelig Software Package

To cite Zelig as a whole, please reference these two sources:

Kosuke Imai, Gary King, and Olivia Lau. 2007. “Zelig: Everyone’s Statistical Software,” <http://GKing.harvard.edu/zelig>.

Imai, Kosuke, Gary King, and Olivia Lau. (2008). “Toward A Common Framework for Statistical Analysis and Development.” *Journal of Computational and Graphical Statistics*, Vol. 17, No. 4 (December), pp. 892-913.

See also

The negative binomial model is part of the MASS package by William N. Venable and Brian D. Ripley [6]. Advanced users may wish to refer to `help(glm.nb)` as well as [5]. Robust standard errors are implemented via sandwich package by Achim Zeileis [7]. Sample data are from [4].

References

- [1] Donald W.K. Andrews. Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica*, 59(3):817–858, May 1991.
- [2] Kosuke Imai, Olivia Lau, and Gary King. *negbinom: Negative Binomial Regression for Event Count Dependent Variables*, 2011.
- [3] Thomas Lumley and Patrick Heagerty. Weighted empirical adaptive variance estimators for correlated data regression. *jrssb*, 61(2):459–477, 1999.

- [4] Lisa Martin. *Coercive Cooperation: Explaining Multilateral Economic Sanctions*. Princeton University Press, 1992. Please inquire with Lisa Martin before publishing results from these data, as this dataset includes errors that have since been corrected.
- [5] Peter McCullagh and James A. Nelder. *Generalized Linear Models*. Number 37 in Monograph on Statistics and Applied Probability. Chapman & Hall, 2nd edition, 1989.
- [6] William N. Venables and Brian D. Ripley. *Modern Applied Statistics with S*. Springer-Verlag, 4th edition, 2002.
- [7] Achim Zeileis. Econometric computing with hc and hac covariance matrix estimators. *Journal of Statistical Software*, 11(10):1–17, 2004.